

LOC169505 AS MODIFIER OF THE APC AND AXIN PATHWAYS AND METHODS OF USE

REFERENCE TO RELATED APPLICATIONS

5 This application claims priority to U.S. provisional patent application 60/429,060 filed 11/25/2002. The contents of the prior applications are hereby incorporated in their entirety.

BACKGROUND OF THE INVENTION

10 Deregulation of beta-catenin signaling is a frequent and early event in the development of a variety of human tumors, including colon cancer, melanoma, ovarian cancer, and prostate cancer. Activation of beta-catenin signaling can occur in tumor cells by loss-of-function mutations in the tumor suppressor genes APC (adenomatus polyposis coli protein) or Axin, as well as by gain-of-function mutations in the oncogene beta-
15 catenin itself. APC, Axin, and beta-catenin normally bind to each other, as well as to the serine/threonine kinase GSK3-beta. Assembly of this degradation complex allows GSK3-beta to phosphorylate beta-catenin, which leads to beta-catenin ubiquitination and degradation by the proteasome. In the absence of APC or Axin activity, beta-catenin protein becomes stabilized and accumulates in the nucleus where it acts as a
20 transcriptional co-activator with TCF for the induction of target genes, including the cell cycle regulators cyclin D1 and c-Myc.

 The *C. elegans* gene *pry-1* is the structural and functional ortholog of vertebrate Axin (Korswagen HC et al. (2002) Genes Dev. 16:1291-302). PRY-1 is predicted to contain conserved RGS and DIX domains that, in Axin, bind APC and Dishevelled,
25 respectively. Overexpression of the *C. elegans pry-1* gene in zebrafish can fully rescue the mutant phenotype of *masterblind*, the zebrafish Axin1 mutation. *pry-1* loss-of-function mutations produce several phenotypes that appear to result from increased beta-catenin signaling (Gleason JE et al. (2002) Genes Dev. 16:1281-90; Korswagen et al., *supra*). We find that the temperature-sensitive, reduction-of-function *pry-1* mutant *mu38*
30 grown at 25°C produces a multivulva (Muv) phenotype in which approximately 30% of animals are induced to form ectopic vulvae. The *pry-1* Muv mutant phenotype is suppressed by RNAi-mediated inactivation the beta-catenin ortholog *bar-1* and the TCF ortholog *pop-1*. The Muv phenotype can also be generated by gain-of-function mutations

in *bar-1*/beta-catenin that eliminate the consensus GSK3-beta phosphorylation sites and are predicted to prevent Axin-mediated degradation of BAR-1.

The *C. elegans* gene product APR-1 shows significant structural similarity to human APC and can bind to both BAR-1/beta-catenin and PRY-1/Axin (Rocheleau et al. (1997), Cell, Vol. 90, 707-716; Natarajan et al. (2001), Genetics, Vol. 159, 159-172; Korswagen et al., *supra*).

The ability to manipulate the genomes of model organisms such as *C. elegans* provides a powerful means to analyze biochemical processes that, due to significant evolutionary conservation, have direct relevance to more complex vertebrate organisms. Due to a high level of gene and pathway conservation, the strong similarity of cellular processes, and the functional conservation of genes between these model organisms and mammals, identification of the involvement of novel genes in particular pathways and their functions in such model organisms can directly contribute to the understanding of the correlative pathways and methods of modulating them in mammals (see, for example, Dulubova I, et al, J Neurochem 2001 Apr;77(1):229-38; Cai T, et al., Diabetologia 2001 Jan;44(1):81-8; Pasquinelli AE, et al., Nature. 2000 Nov 2;408(6808):37-8; Ivanov IP, et al., EMBO J 2000 Apr 17;19(8):1907-17; Vajo Z et al., Mamm Genome 1999 Oct;10(10):1000-4). For example, a genetic screen can be carried out in an invertebrate model organism having underexpression (e.g. knockout) or overexpression of a gene (referred to as a "genetic entry point") that yields a visible phenotype. Additional genes are mutated in a random or targeted manner. When a gene mutation changes the original phenotype caused by the mutation in the genetic entry point, the gene is identified as a "modifier" involved in the same or overlapping pathway as the genetic entry point. When the genetic entry point is an ortholog of a human gene implicated in a disease pathway, such as APC or axin, modifier genes can be identified that may be attractive candidate targets for novel therapeutics.

All references cited herein, including patents, patent applications, publications, and sequence information in referenced Genbank identifier numbers, are incorporated herein in their entireties.

SUMMARY OF THE INVENTION

We have discovered genes that modify the APC and axin pathways in *C. elegans*, and identified their human orthologs, hereinafter referred to as LOC169505. The invention provides methods for utilizing these APC and axin modifier genes and

polypeptides to identify LOC169505-modulating agents that are candidate therapeutic agents that can be used in the treatment of disorders associated with defective or impaired APC or axin function and/or LOC169505 function. Preferred LOC169505-modulating agents specifically bind to LOC169505 polypeptides and restore APC or axin function.

- 5 Other preferred LOC169505-modulating agents are nucleic acid modulators such as antisense oligomers and RNAi that repress LOC169505 gene expression or product activity by, for example, binding to and inhibiting the respective nucleic acid (i.e. DNA or mRNA).

- LOC169505 modulating agents may be evaluated by any convenient *in vitro* or *in vivo* assay for molecular interaction with an LOC169505 polypeptide or nucleic acid. In one embodiment, candidate LOC169505 modulating agents are tested with an assay system comprising a LOC169505 polypeptide or nucleic acid. Agents that produce a change in the activity of the assay system relative to controls are identified as candidate APC and axin modulating agents. The assay system may be cell-based or cell-free.
- 15 LOC169505-modulating agents include LOC169505 related proteins (e.g. dominant negative mutants, and biotherapeutics); LOC169505 -specific antibodies; LOC169505 -specific antisense oligomers and other nucleic acid modulators; and chemical agents that specifically bind to or interact with LOC169505 or compete with LOC169505 binding partner (e.g. by binding to an LOC169505 binding partner). In one specific embodiment,
- 20 a small molecule modulator is identified using a kinase assay. In specific embodiments, the screening assay system is selected from a binding assay, an apoptosis assay, a cell proliferation assay, an angiogenesis assay, and a hypoxic induction assay.

- In another embodiment, candidate APC and axin pathway modulating agents are further tested using a second assay system that detects changes in the APC and axin pathways, such as angiogenic, apoptotic, or cell proliferation changes produced by the originally identified candidate agent or an agent derived from the original agent. The second assay system may use cultured cells or non-human animals. In specific embodiments, the secondary assay system uses non-human animals, including animals predetermined to have a disease or disorder implicating the APC and axin pathways, such as an angiogenic, apoptotic, or cell proliferation disorder (e.g. cancer).
- 25
- 30

The invention further provides methods for modulating the LOC169505 function and/or the APC and axin pathways in a mammalian cell by contacting the mammalian cell with an agent that specifically binds a LOC169505 polypeptide or nucleic acid. The agent may be a small molecule modulator, a nucleic acid modulator, or an antibody and may be

administered to a mammalian animal predetermined to have a pathology associated with the APC and axin pathways.

DETAILED DESCRIPTION OF THE INVENTION

5 Genetic screens were designed to identify modifiers of the axin and APC pathway in *C. elegans*. The function of *ap-1* was depleted by RNAi in a *pry-1* hypomorphic allele *mu38*. Various specific genes were then silenced by RNA inhibition (RNAi). Methods for using RNAi to silence genes in *C. elegans* are known in the art (Fire A, et al., 1998 Nature 391:806-811; Fire, A. Trends Genet. 15, 358-363 (1999); WO9932619). Genes
10 causing altered phenotypes in the worms were identified as modifiers of the APC and axin pathways. The F26D11.1 gene was identified as a modifier of the APC and axin pathways. Accordingly, vertebrate orthologs of this modifier, and preferably the human orthologs, LOC169505 genes (i.e., nucleic acids and polypeptides) are attractive drug targets for the treatment of pathologies associated with a defective APC and axin signaling
15 pathway, such as cancer.

In vitro and in vivo methods of assessing LOC169505 function are provided herein. Modulation of the LOC169505 or their respective binding partners is useful for understanding the association of the APC and axin pathways and their members in normal and disease conditions and for developing diagnostics and therapeutic modalities for APC
20 and axin related pathologies. LOC169505-modulating agents that act by inhibiting or enhancing LOC169505 expression, directly or indirectly, for example, by affecting an LOC169505 function such as enzymatic (e.g., catalytic) or binding activity, can be identified using methods provided herein. LOC169505 modulating agents are useful in diagnosis, therapy and pharmaceutical development.

25

Nucleic acids and polypeptides of the invention

Sequences related to LOC169505 nucleic acids and polypeptides that can be used in the invention are disclosed in Genbank (referenced by Genbank identifier (GI) number as GI#s 20537768 (SEQ ID NO:1) and 29731938 (SEQ ID NO:2) for nucleic acid, and
30 GI# 20537769 (SEQ ID NO:5) for polypeptide. Additionally, nucleic acid sequences of SEQ ID NOs:3 and 4 can also be used in the invention.

The term "LOC169505 polypeptide" refers to a full-length LOC169505 protein or a functionally active fragment or derivative thereof. A "functionally active" LOC169505 fragment or derivative exhibits one or more functional activities associated with a full-

length, wild-type LOC169505 protein, such as antigenic or immunogenic activity, enzymatic activity, ability to bind natural cellular substrates, etc. The functional activity of LOC169505 proteins, derivatives and fragments can be assayed by various methods known to one skilled in the art (Current Protocols in Protein Science (1998) Coligan *et al.*,
5 eds., John Wiley & Sons, Inc., Somerset, New Jersey) and as further discussed below. In one embodiment, a functionally active LOC169505 polypeptide is a LOC169505 derivative capable of rescuing defective endogenous LOC169505 activity, such as in cell based or animal assays; the rescuing derivative may be from the same or a different species. For purposes herein, functionally active fragments also include those fragments
10 that comprise one or more structural domains of an LOC169505, such as a kinase domain or a binding domain. Protein domains can be identified using the PFAM program (Bateman A., et al., Nucleic Acids Res, 1999, 27:260-2). For example, the kinase domain of LOC169505 from GI# 20537769 (SEQ ID NO:5) is located at approximately amino acid residues 7 to 186. Methods for obtaining LOC169505 polypeptides are also further
15 described below. In some embodiments, preferred fragments are functionally active, domain-containing fragments comprising at least 25 contiguous amino acids, preferably at least 50, more preferably 75, and most preferably at least 100 contiguous amino acids of an LOC169505. In further preferred embodiments, the fragment comprises the entire kinase (functionally active) domain.

20 The term "LOC169505 nucleic acid" refers to a DNA or RNA molecule that encodes a LOC169505 polypeptide. Preferably, the LOC169505 polypeptide or nucleic acid or fragment thereof is from a human, but can also be an ortholog, or derivative thereof with at least 70% sequence identity, preferably at least 80%, more preferably 85%, still more preferably 90%, and most preferably at least 95% sequence identity with human
25 LOC169505. Methods of identifying orthologs are known in the art. Normally, orthologs in different species retain the same function, due to presence of one or more protein motifs and/or 3-dimensional structures. Orthologs are generally identified by sequence homology analysis, such as BLAST analysis, usually using protein bait sequences. Sequences are assigned as a potential ortholog if the best hit sequence from the forward BLAST result
30 retrieves the original query sequence in the reverse BLAST (Huynen MA and Bork P, Proc Natl Acad Sci (1998) 95:5849-5856; Huynen MA *et al.*, Genome Research (2000) 10:1204-1210). Programs for multiple sequence alignment, such as CLUSTAL (Thompson JD et al, 1994, Nucleic Acids Res 22:4673-4680) may be used to highlight conserved regions and/or residues of orthologous proteins and to generate phylogenetic

trees. In a phylogenetic tree representing multiple homologous sequences from diverse species (e.g., retrieved through BLAST analysis), orthologous sequences from two species generally appear closest on the tree with respect to all other sequences from these two species. Structural threading or other analysis of protein folding (e.g., using software by ProCeryon, Biosciences, Salzburg, Austria) may also identify potential orthologs. In evolution, when a gene duplication event follows speciation, a single gene in one species, such as *C.elegans*, may correspond to multiple genes (paralogs) in another, such as human. As used herein, the term "orthologs" encompasses paralogs. As used herein, "percent (%) sequence identity" with respect to a subject sequence, or a specified portion of a subject sequence, is defined as the percentage of nucleotides or amino acids in the candidate derivative sequence identical with the nucleotides or amino acids in the subject sequence (or specified portion thereof), after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent sequence identity, as generated by the program WU-BLAST-2.0a19 (Altschul *et al.*, J. Mol. Biol. (1997) 215:403-410) with all the search parameters set to default values. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched. A % identity value is determined by the number of matching identical nucleotides or amino acids divided by the sequence length for which the percent identity is being reported. "Percent (%) amino acid sequence similarity" is determined by doing the same calculation as for determining % amino acid sequence identity, but including conservative amino acid substitutions in addition to identical amino acids in the computation.

A conservative amino acid substitution is one in which an amino acid is substituted for another amino acid having similar properties such that the folding or activity of the protein is not significantly affected. Aromatic amino acids that can be substituted for each other are phenylalanine, tryptophan, and tyrosine; interchangeable hydrophobic amino acids are leucine, isoleucine, methionine, and valine; interchangeable polar amino acids are glutamine and asparagine; interchangeable basic amino acids are arginine, lysine and histidine; interchangeable acidic amino acids are aspartic acid and glutamic acid; and interchangeable small amino acids are alanine, serine, threonine, cysteine and glycine.

Alternatively, an alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman (Smith and Waterman, 1981, Advances in Applied Mathematics 2:482-489; database: European Bioinformatics Institute; Smith and

Waterman, 1981, J. of Molec.Biol., 147:195-197; Nicholas et al., 1998, "A Tutorial on Searching Sequence Databases and Sequence Scoring Methods" (www.psc.edu) and references cited therein.; W.R. Pearson, 1991, Genomics 11:635-650). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff (Dayhoff: Atlas of Protein Sequences and Structure, M. O. Dayhoff ed., 5 suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA), and normalized by Gribskov (Gribskov 1986 Nucl. Acids Res. 14(6):6745-6763). The Smith-Waterman algorithm may be employed where default parameters are used for scoring (for example, gap open penalty of 12, gap extension penalty of two). From the data generated, the "Match" value reflects "sequence identity."

Derivative nucleic acid molecules of the subject nucleic acid molecules include sequences that hybridize to the nucleic acid sequence of an LOC169505. The stringency of hybridization can be controlled by temperature, ionic strength, pH, and the presence of denaturing agents such as formamide during hybridization and washing. Conditions routinely used are set out in readily available procedure texts (*e.g.*, Current Protocol in Molecular Biology, Vol. 1, Chap. 2.10, John Wiley & Sons, Publishers (1994); Sambrook *et al.*, Molecular Cloning, Cold Spring Harbor (1989)). In some embodiments, a nucleic acid molecule of the invention is capable of hybridizing to a nucleic acid molecule containing the nucleotide sequence of an LOC169505 under high stringency hybridization conditions that are: prehybridization of filters containing nucleic acid for 8 hours to overnight at 65° C in a solution comprising 6X single strength citrate (SSC) (1X SSC is 0.15 M NaCl, 0.015 M Na citrate; pH 7.0), 5X Denhardt's solution, 0.05% sodium pyrophosphate and 100 µg/ml herring sperm DNA; hybridization for 18-20 hours at 65° C in a solution containing 6X SSC, 1X Denhardt's solution, 100 µg/ml yeast tRNA and 0.05% sodium pyrophosphate; and washing of filters at 65° C for 1h in a solution containing 0.1X SSC and 0.1% SDS (sodium dodecyl sulfate).

In other embodiments, moderately stringent hybridization conditions are used that are: pretreatment of filters containing nucleic acid for 6 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA; hybridization for 18-20h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, and 10% (wt/vol) dextran sulfate; followed by washing twice for 1 hour at 55° C in a solution containing 2X SSC and 0.1% SDS.

Alternatively, low stringency conditions can be used that are: incubation for 8 hours to overnight at 37° C in a solution comprising 20% formamide, 5 x SSC, 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured sheared salmon sperm DNA; hybridization in the same buffer for 18 to 20
5 hours; and washing of filters in 1 x SSC at about 37° C for 1 hour.

Isolation, Production, Expression, and Mis-expression of LOC169505 Nucleic Acids and Polypeptides

LOC169505 nucleic acids and polypeptides are useful for identifying and testing
10 agents that modulate LOC169505 function and for other applications related to the involvement of LOC169505 in the APC and axin pathways. LOC169505 nucleic acids and derivatives and orthologs thereof may be obtained using any available method. For instance, techniques for isolating cDNA or genomic DNA sequences of interest by screening DNA libraries or by using polymerase chain reaction (PCR) are well known in
15 the art. In general, the particular use for the protein will dictate the particulars of expression, production, and purification methods. For instance, production of proteins for use in screening for modulating agents may require methods that preserve specific biological activities of these proteins, whereas production of proteins for antibody generation may require structural integrity of particular epitopes. Expression of proteins
20 to be purified for screening or antibody production may require the addition of specific tags (*e.g.*, generation of fusion proteins). Overexpression of an LOC169505 protein for assays used to assess LOC169505 function, such as involvement in cell cycle regulation or hypoxic response, may require expression in eukaryotic cell lines capable of these cellular activities. Techniques for the expression, production, and purification of proteins are well
25 known in the art; any suitable means therefore may be used (*e.g.*, Higgins SJ and Hames BD (eds.) Protein Expression: A Practical Approach, Oxford University Press Inc., New York 1999; Stanbury PF et al., Principles of Fermentation Technology, 2nd edition, Elsevier Science, New York, 1995; Doonan S (ed.) Protein Purification Protocols, Humana Press, New Jersey, 1996; Coligan JE et al, Current Protocols in Protein Science
30 (eds.), 1999, John Wiley & Sons, New York). In particular embodiments, recombinant LOC169505 is expressed in a cell line known to have defective APC and axin function. The recombinant cells are used in cell-based screening assay systems of the invention, as described further below.

The nucleotide sequence encoding an LOC169505 polypeptide can be inserted into any appropriate expression vector. The necessary transcriptional and translational signals, including promoter/enhancer element, can derive from the native LOC169505 gene and/or its flanking regions or can be heterologous. A variety of host-vector expression systems
5 may be utilized, such as mammalian cell systems infected with virus (*e.g.* vaccinia virus, adenovirus, *etc.*); insect cell systems infected with virus (*e.g.* baculovirus); microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage, plasmid, or cosmid DNA. An isolated host cell strain that modulates the expression of, modifies, and/or specifically processes the gene product may be used.

10 To detect expression of the LOC169505 gene product, the expression vector can comprise a promoter operably linked to an LOC169505 gene nucleic acid, one or more origins of replication, and, one or more selectable markers (*e.g.* thymidine kinase activity, resistance to antibiotics, *etc.*). Alternatively, recombinant expression vectors can be identified by assaying for the expression of the LOC169505 gene product based on the
15 physical or functional properties of the LOC169505 protein in *in vitro* assay systems (*e.g.* immunoassays).

The LOC169505 protein, fragment, or derivative may be optionally expressed as a fusion, or chimeric protein product (i.e. it is joined via a peptide bond to a heterologous protein sequence of a different protein), for example to facilitate purification or detection.

20 A chimeric product can be made by ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences to each other using standard methods and expressing the chimeric product. A chimeric product may also be made by protein synthetic techniques, *e.g.* by use of a peptide synthesizer (Hunkapiller *et al.*, Nature (1984) 310:105-111).

25 Once a recombinant cell that expresses the LOC169505 gene sequence is identified, the gene product can be isolated and purified using standard methods (*e.g.* ion exchange, affinity, and gel exclusion chromatography; centrifugation; differential solubility; electrophoresis). Alternatively, native LOC169505 proteins can be purified from natural sources, by standard methods (*e.g.* immunoaffinity purification). Once a
30 protein is obtained, it may be quantified and its activity measured by appropriate methods, such as immunoassay, bioassay, or other measurements of physical properties, such as crystallography.

The methods of this invention may also use cells that have been engineered for altered expression (mis-expression) of LOC169505 or other genes associated with the

APC and axin pathways. As used herein, mis-expression encompasses ectopic expression, over-expression, under-expression, and non-expression (*e.g.* by gene knock-out or blocking expression that would otherwise normally occur).

5 Genetically modified animals

Animal models that have been genetically modified to alter LOC169505 expression may be used in *in vivo* assays to test for activity of a candidate APC and axin modulating agent, or to further assess the role of LOC169505 in a APC and axin pathways process such as apoptosis or cell proliferation. Preferably, the altered LOC169505 expression results in a detectable phenotype, such as decreased or increased levels of cell proliferation, angiogenesis, or apoptosis compared to control animals having normal LOC169505 expression. The genetically modified animal may additionally have altered APC or axin expression (*e.g.* APC or axin knockout). Preferred genetically modified animals are mammals such as primates, rodents (preferably mice or rats), among others. Preferred non-mammalian species include zebrafish, *C. elegans*, and *Drosophila*. Preferred genetically modified animals are transgenic animals having a heterologous nucleic acid sequence present as an extrachromosomal element in a portion of its cells, *i.e.* mosaic animals (see, for example, techniques described by Jakobovits, 1994, Curr. Biol. 4:761-763.) or stably integrated into its germ line DNA (*i.e.*, in the genomic sequence of most or all of its cells). Heterologous nucleic acid is introduced into the germ line of such transgenic animals by genetic manipulation of, for example, embryos or embryonic stem cells of the host animal.

Methods of making transgenic animals are well-known in the art (for transgenic mice see Brinster *et al.*, Proc. Nat. Acad. Sci. USA 82: 4438-4442 (1985), U.S. Pat. Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Pat. No. 4,873,191 by Wagner *et al.*, and Hogan, B., Manipulating the Mouse Embryo, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1986); for particle bombardment see U.S. Pat. No., 4,945,050, by Sandford *et al.*; for transgenic *Drosophila* see Rubin and Spradling, Science (1982) 218:348-53 and U.S. Pat. No. 4,670,388; for transgenic insects see Berghammer A.J. *et al.*, A Universal Marker for Transgenic Insects (1999) Nature 402:370-371; for transgenic Zebrafish see Lin S., Transgenic Zebrafish, Methods Mol Biol. (2000);136:375-3830; for microinjection procedures for fish, amphibian eggs and birds see Houdebine and Chourrout, Experientia (1991) 47:897-905; for transgenic rats see Hammer *et al.*, Cell (1990) 63:1099-1112; and for culturing of embryonic stem (ES) cells and the subsequent

production of transgenic animals by the introduction of DNA into ES cells using methods such as electroporation, calcium phosphate/DNA precipitation and direct injection see, e.g., *Teratocarcinomas and Embryonic Stem Cells, A Practical Approach*, E. J. Robertson, ed., IRL Press (1987)). Clones of the nonhuman transgenic animals can be produced
5 according to available methods (see Wilmut, I. *et al.* (1997) *Nature* 385:810-813; and PCT International Publication Nos. WO 97/07668 and WO 97/07669).

In one embodiment, the transgenic animal is a "knock-out" animal having a heterozygous or homozygous alteration in the sequence of an endogenous LOC169505 gene that results in a decrease of LOC169505 function, preferably such that LOC169505
10 expression is undetectable or insignificant. Knock-out animals are typically generated by homologous recombination with a vector comprising a transgene having at least a portion of the gene to be knocked out. Typically a deletion, addition or substitution has been introduced into the transgene to functionally disrupt it. The transgene can be a human gene (e.g., from a human genomic clone) but more preferably is an ortholog of the human
15 gene derived from the transgenic host species. For example, a mouse LOC169505 gene is used to construct a homologous recombination vector suitable for altering an endogenous LOC169505 gene in the mouse genome. Detailed methodologies for homologous recombination in mice are available (see Capecchi, *Science* (1989) 244:1288-1292; Joyner *et al.*, *Nature* (1989) 338:153-156). Procedures for the production of non-rodent
20 transgenic mammals and other animals are also available (Houdebine and Chourrout, *supra*; Pursel *et al.*, *Science* (1989) 244:1281-1288; Simms *et al.*, *Bio/Technology* (1988) 6:179-183). In a preferred embodiment, knock-out animals, such as mice harboring a knockout of a specific gene, may be used to produce antibodies against the human counterpart of the gene that has been knocked out (Claesson MH *et al.*, (1994) *Scan J*
25 *Immunol* 40:257-264; Declerck PJ *et al.*, (1995) *J Biol Chem.* 270:8397-400).

In another embodiment, the transgenic animal is a "knock-in" animal having an alteration in its genome that results in altered expression (e.g., increased (including ectopic) or decreased expression) of the LOC169505 gene, e.g., by introduction of additional copies of LOC169505, or by operatively inserting a regulatory sequence that
30 provides for altered expression of an endogenous copy of the LOC169505 gene. Such regulatory sequences include inducible, tissue-specific, and constitutive promoters and enhancer elements. The knock-in can be homozygous or heterozygous.

Transgenic nonhuman animals can also be produced that contain selected systems allowing for regulated expression of the transgene. One example of such a system that

may be produced is the cre/loxP recombinase system of bacteriophage P1 (Lakso *et al.*, PNAS (1992) 89:6232-6236; U.S. Pat. No. 4,959,317). If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be
5 provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman *et al.* (1991) Science 251:1351-1355; U.S. Pat. No. 5,654,182). In a preferred embodiment,
10 both Cre-LoxP and Flp-Frt are used in the same system to regulate expression of the transgene, and for sequential deletion of vector sequences in the same cell (Sun X *et al.* (2000) Nat Genet 25:83-6).

The genetically modified animals can be used in genetic studies to further elucidate the APC and axin pathways, as animal models of disease and disorders implicating
15 defective APC and axin function, and for *in vivo* testing of candidate therapeutic agents, such as those identified in screens described below. The candidate therapeutic agents are administered to a genetically modified animal having altered LOC169505 function and phenotypic changes are compared with appropriate control animals such as genetically modified animals that receive placebo treatment, and/or animals with unaltered
20 LOC169505 expression that receive candidate therapeutic agent.

In addition to the above-described genetically modified animals having altered LOC169505 function, animal models having defective APC and axin function (and otherwise normal LOC169505 function), can be used in the methods of the present invention. For example, a APC or axin knockout mouse can be used to assess, *in vivo*, the
25 activity of a candidate APC and axin modulating agent identified in one of the *in vitro* assays described below. Preferably, the candidate APC and axin modulating agent when administered to a model system with cells defective in APC and axin function, produces a detectable phenotypic change in the model system indicating that the APC and axin function is restored, i.e., the cells exhibit normal cell cycle progression.

30

Modulating Agents

The invention provides methods to identify agents that interact with and/or modulate the function of LOC169505 and/or the APC and axin pathways. Modulating agents identified by the methods are also part of the invention. Such agents are useful in a

variety of diagnostic and therapeutic applications associated with the APC and axin pathways, as well as in further analysis of the LOC169505 protein and its contribution to the APC and axin pathways. Accordingly, the invention also provides methods for modulating the APC and axin pathways comprising the step of specifically modulating
5 LOC169505 activity by administering a LOC169505-interacting or -modulating agent.

As used herein, an "LOC169505-modulating agent" is any agent that modulates LOC169505 function, for example, an agent that interacts with LOC169505 to inhibit or enhance LOC169505 activity or otherwise affect normal LOC169505 function. LOC169505 function can be affected at any level, including transcription, protein
10 expression, protein localization, and cellular or extra-cellular activity. In a preferred embodiment, the LOC169505 - modulating agent specifically modulates the function of the LOC169505. The phrases "specific modulating agent", "specifically modulates", etc., are used herein to refer to modulating agents that directly bind to the LOC169505 polypeptide or nucleic acid, and preferably inhibit, enhance, or otherwise alter, the
15 function of the LOC169505. These phrases also encompass modulating agents that alter the interaction of the LOC169505 with a binding partner, substrate, or cofactor (e.g. by binding to a binding partner of an LOC169505, or to a protein/binding partner complex, and altering LOC169505 function). In a further preferred embodiment, the LOC169505-modulating agent is a modulator of the APC and axin pathways (e.g. it restores and/or
20 upregulates APC and axin function) and thus is also a APC and axin-modulating agent.

Preferred LOC169505-modulating agents include small molecule compounds; LOC169505-interacting proteins, including antibodies and other biotherapeutics; and nucleic acid modulators such as antisense and RNA inhibitors. The modulating agents may be formulated in pharmaceutical compositions, for example, as compositions that
25 may comprise other active ingredients, as in combination therapy, and/or suitable carriers or excipients. Techniques for formulation and administration of the compounds may be found in "Remington's Pharmaceutical Sciences" Mack Publishing Co., Easton, PA, 19th edition.

30 **Small molecule modulators**

Small molecules are often preferred to modulate function of proteins with enzymatic function, and/or containing protein interaction domains. Chemical agents, referred to in the art as "small molecule" compounds are typically organic, non-peptide molecules, having a molecular weight up to 10,000, preferably up to 5,000, more

preferably up to 1,000, and most preferably up to 500 daltons. This class of modulators includes chemically synthesized molecules, for instance, compounds from combinatorial chemical libraries. Synthetic compounds may be rationally designed or identified based on known or inferred properties of the LOC169505 protein or may be identified by
5 screening compound libraries. Alternative appropriate modulators of this class are natural products, particularly secondary metabolites from organisms such as plants or fungi, which can also be identified by screening compound libraries for LOC169505-modulating activity. Methods for generating and obtaining compounds are well known in the art (Schreiber SL, Science (2000) 151: 1964-1969; Radmann J and Gunther J, Science (2000)
10 151:1947-1948).

Small molecule modulators identified from screening assays, as described below, can be used as lead compounds from which candidate clinical compounds may be designed, optimized, and synthesized. Such clinical compounds may have utility in treating pathologies associated with the APC and axin pathways. The activity of candidate
15 small molecule modulating agents may be improved several-fold through iterative secondary functional validation, as further described below, structure determination, and candidate modulator modification and testing. Additionally, candidate clinical compounds are generated with specific regard to clinical and pharmacological properties. For example, the reagents may be derivatized and re-screened using *in vitro* and *in vivo* assays to
20 optimize activity and minimize toxicity for pharmaceutical development.

Protein Modulators

Specific LOC169505-interacting proteins are useful in a variety of diagnostic and therapeutic applications related to the APC and axin pathways and related disorders, as
25 well as in validation assays for other LOC169505-modulating agents. In a preferred embodiment, LOC169505-interacting proteins affect normal LOC169505 function, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In another embodiment, LOC169505-interacting proteins are useful in detecting and providing information about the function of LOC169505 proteins, as is
30 relevant to APC and axin related disorders, such as cancer (e.g., for diagnostic means).

An LOC169505-interacting protein may be endogenous, i.e. one that naturally interacts genetically or biochemically with an LOC169505, such as a member of the LOC169505 pathway that modulates LOC169505 expression, localization, and/or activity. LOC169505-modulators include dominant negative forms of LOC169505-interacting

proteins and of LOC169505 proteins themselves. Yeast two-hybrid and variant screens offer preferred methods for identifying endogenous LOC169505-interacting proteins (Finley, R. L. et al. (1996) in DNA Cloning-Expression Systems: A Practical Approach, eds. Glover D. & Hames B. D (Oxford University Press, Oxford, England), pp. 169-203; 5 Fashema SF et al., Gene (2000) 250:1-14; Drees BL Curr Opin Chem Biol (1999) 3:64-70; Vidal M and Legrain P Nucleic Acids Res (1999) 27:919-29; and U.S. Pat. No. 5,928,868). Mass spectrometry is an alternative preferred method for the elucidation of protein complexes (reviewed in, e.g., Pandley A and Mann M, Nature (2000) 405:837-846; Yates JR 3rd, Trends Genet (2000) 16:5-8).

10 An LOC169505-interacting protein may be an exogenous protein, such as an LOC169505-specific antibody or a T-cell antigen receptor (see, e.g., Harlow and Lane (1988) Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory; Harlow and Lane (1999) Using antibodies: a laboratory manual. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press). LOC169505 antibodies are further discussed below.

15 In preferred embodiments, an LOC169505-interacting protein specifically binds an LOC169505 protein. In alternative preferred embodiments, an LOC169505-modulating agent binds an LOC169505 substrate, binding partner, or cofactor.

Antibodies

20 In another embodiment, the protein modulator is an LOC169505 specific antibody agonist or antagonist. The antibodies have therapeutic and diagnostic utilities, and can be used in screening assays to identify LOC169505 modulators. The antibodies can also be used in dissecting the portions of the LOC169505 pathway responsible for various cellular responses and in the general processing and maturation of the LOC169505.

25 Antibodies that specifically bind LOC169505 polypeptides can be generated using known methods. Preferably the antibody is specific to a mammalian ortholog of LOC169505 polypeptide, and more preferably, to human LOC169505. Antibodies may be polyclonal, monoclonal (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a FAb
30 expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. Epitopes of LOC169505 which are particularly antigenic can be selected, for example, by routine screening of LOC169505 polypeptides for antigenicity or by applying a theoretical method for selecting antigenic regions of a protein (Hopp and Wood (1981), Proc. Nati. Acad. Sci. U.S.A. 78:3824-28; Hopp and Wood, (1983) Mol.

Immunol. 20:483-89; Sutcliffe et al., (1983) Science 219:660-66) to the amino acid sequence of an LOC169505. Monoclonal antibodies with affinities of 10^8 M^{-1} preferably 10^9 M^{-1} to 10^{10} M^{-1} , or stronger can be made by standard procedures as described (Harlow and Lane, *supra*; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed) Academic Press, New York; and U.S. Pat. Nos. 4,381,292; 4,451,570; and 4,618,577).
5 Antibodies may be generated against crude cell extracts of LOC169505 or substantially purified fragments thereof. If LOC169505 fragments are used, they preferably comprise at least 10, and more preferably, at least 20 contiguous amino acids of an LOC169505 protein. In a particular embodiment, LOC169505-specific antigens and/or immunogens
10 are coupled to carrier proteins that stimulate the immune response. For example, the subject polypeptides are covalently coupled to the keyhole limpet hemocyanin (KLH) carrier, and the conjugate is emulsified in Freund's complete adjuvant, which enhances the immune response. An appropriate immune system such as a laboratory rabbit or mouse is immunized according to conventional protocols.

15 The presence of LOC169505-specific antibodies is assayed by an appropriate assay such as a solid phase enzyme-linked immunosorbant assay (ELISA) using immobilized corresponding LOC169505 polypeptides. Other assays, such as radioimmunoassays or fluorescent assays might also be used.

Chimeric antibodies specific to LOC169505 polypeptides can be made that contain
20 different portions from different animal species. For instance, a human immunoglobulin constant region may be linked to a variable region of a murine mAb, such that the antibody derives its biological activity from the human antibody, and its binding specificity from the murine fragment. Chimeric antibodies are produced by splicing together genes that encode the appropriate regions from each species (Morrison et al.,
25 Proc. Natl. Acad. Sci. (1984) 81:6851-6855; Neuberger et al., Nature (1984) 312:604-608; Takeda et al., Nature (1985) 31:452-454). Humanized antibodies, which are a form of chimeric antibodies, can be generated by grafting complementary-determining regions (CDRs) (Carlos, T. M., J. M. Harlan. 1994. Blood 84:2068-2101) of mouse antibodies into a background of human framework regions and constant regions by recombinant
30 DNA technology (Riechmann LM, et al., 1988 Nature 323: 323-327). Humanized antibodies contain ~10% murine sequences and ~90% human sequences, and thus further reduce or eliminate immunogenicity, while retaining the antibody specificities (Co MS, and Queen C. 1991 Nature 351: 501-501; Morrison SL. 1992 Ann. Rev. Immun.

10:239-265). Humanized antibodies and methods of their production are well-known in the art (U.S. Pat. Nos. 5,530,101, 5,585,089, 5,693,762, and 6,180,370).

LOC169505-specific single chain antibodies which are recombinant, single chain polypeptides formed by linking the heavy and light chain fragments of the Fv regions via
5 an amino acid bridge, can be produced by methods known in the art (U.S. Pat. No. 4,946,778; Bird, Science (1988) 242:423-426; Huston et al., Proc. Natl. Acad. Sci. USA (1988) 85:5879-5883; and Ward et al., Nature (1989) 334:544-546).

Other suitable techniques for antibody production involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of
10 antibodies in phage or similar vectors (Huse et al., Science (1989) 246:1275-1281). As used herein, T-cell antigen receptors are included within the scope of antibody modulators (Harlow and Lane, 1988, *supra*).

The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, antibodies will be labeled by joining, either covalently
15 or non-covalently, a substance that provides for a detectable signal, or that is toxic to cells that express the targeted protein (Menard S, et al., Int J. Biol Markers (1989) 4:131-134). A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include
20 radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, fluorescent emitting lanthanide metals, chemiluminescent moieties, bioluminescent moieties, magnetic particles, and the like (U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241). Also, recombinant immunoglobulins may be produced (U.S. Pat. No. 4,816,567). Antibodies to cytoplasmic polypeptides may be delivered and reach their targets by conjugation with membrane-penetrating toxin
25 proteins (U.S. Pat. No. 6,086,900).

When used therapeutically in a patient, the antibodies of the subject invention are typically administered parenterally, when possible at the target site, or intravenously. The therapeutically effective dose and dosage regimen is determined by clinical studies. Typically, the amount of antibody administered is in the range of about 0.1 mg/kg –to
30 about 10 mg/kg of patient weight. For parenteral administration, the antibodies are formulated in a unit dosage injectable form (e.g., solution, suspension, emulsion) in association with a pharmaceutically acceptable vehicle. Such vehicles are inherently nontoxic and non-therapeutic. Examples are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Nonaqueous vehicles such as fixed oils, ethyl

oleate, or liposome carriers may also be used. The vehicle may contain minor amounts of additives, such as buffers and preservatives, which enhance isotonicity and chemical stability or otherwise enhance therapeutic potential. The antibodies' concentrations in such vehicles are typically in the range of about 1 mg/ml to about 10 mg/ml.

- 5 Immunotherapeutic methods are further described in the literature (US Pat. No. 5,859,206; WO0073469).

Nucleic Acid Modulators

- Other preferred LOC169505-modulating agents comprise nucleic acid molecules,
10 such as antisense oligomers or double stranded RNA (dsRNA), which generally inhibit LOC169505 activity. Preferred nucleic acid modulators interfere with the function of the LOC169505 nucleic acid such as DNA replication, transcription, translocation of the LOC169505 RNA to the site of protein translation, translation of protein from the LOC169505 RNA, splicing of the LOC169505 RNA to yield one or more mRNA species,
15 or catalytic activity which may be engaged in or facilitated by the LOC169505 RNA.

- In one embodiment, the antisense oligomer is an oligonucleotide that is sufficiently complementary to an LOC169505 mRNA to bind to and prevent translation, preferably by binding to the 5' untranslated region. LOC169505-specific antisense oligonucleotides, preferably range from at least 6 to about 200 nucleotides. In some embodiments the
20 oligonucleotide is preferably at least 10, 15, or 20 nucleotides in length. In other embodiments, the oligonucleotide is preferably less than 50, 40, or 30 nucleotides in length. The oligonucleotide can be DNA or RNA or a chimeric mixture or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide
25 may include other appending groups such as peptides, agents that facilitate transport across the cell membrane, hybridization-triggered cleavage agents, and intercalating agents.

- In another embodiment, the antisense oligomer is a phosphothioate morpholino oligomer (PMO). PMOs are assembled from four different morpholino subunits, each of
30 which contain one of four genetic bases (A, C, G, or T) linked to a six-membered morpholine ring. Polymers of these subunits are joined by non-ionic phosphodiarnidate intersubunit linkages. Details of how to make and use PMOs and other antisense oligomers are well known in the art (e.g. see WO99/18193; Probst JC, Antisense Oligodeoxynucleotide and Ribozyme Design, Methods. (2000) 22(3):271-281; Summerton

J, and Weller D. 1997 Antisense Nucleic Acid Drug Dev. :7:187-95; US Pat. No. 5,235,033; and US Pat No. 5,378,841).

Alternative preferred LOC169505 nucleic acid modulators are double-stranded RNA species mediating RNA interference (RNAi). RNAi is the process of sequence-specific, post-transcriptional gene silencing in animals and plants, initiated by double-stranded RNA (dsRNA) that is homologous in sequence to the silenced gene. Methods relating to the use of RNAi to silence genes in *C. elegans*, *Drosophila*, plants, and humans are known in the art (Fire A, et al., 1998 Nature 391:806-811; Fire, A. Trends Genet. 15, 358-363 (1999); Sharp, P. A. RNA interference 2001. Genes Dev. 15, 485-490 (2001); Hammond, S. M., et al., Nature Rev. Genet. 2, 110-1119 (2001); Tuschl, T. Chem. Biochem. 2, 239-245 (2001); Hamilton, A. et al., Science 286, 950-952 (1999); Hammond, S. M., et al., Nature 404, 293-296 (2000); Zamore, P. D., et al., Cell 101, 25-33 (2000); Bernstein, E., et al., Nature 409, 363-366 (2001); Elbashir, S. M., et al., Genes Dev. 15, 188-200 (2001); WO0129058; WO9932619; Elbashir SM, et al., 2001 Nature 411:494-498).

Nucleic acid modulators are commonly used as research reagents, diagnostics, and therapeutics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used to elucidate the function of particular genes (see, for example, U.S. Pat. No. 6,165,790). Nucleic acid modulators are also used, for example, to distinguish between functions of various members of a biological pathway. For example, antisense oligomers have been employed as therapeutic moieties in the treatment of disease states in animals and man and have been demonstrated in numerous clinical trials to be safe and effective (Milligan JF, et al, Current Concepts in Antisense Drug Design, J Med Chem. (1993) 36:1923-1937; Tonkinson JL et al., Antisense Oligodeoxynucleotides as Clinical Therapeutic Agents, Cancer Invest. (1996) 14:54-65). Accordingly, in one aspect of the invention, an LOC169505-specific nucleic acid modulator is used in an assay to further elucidate the role of the LOC169505 in the APC and axin pathways, and/or its relationship to other members of the pathway. In another aspect of the invention, an LOC169505-specific antisense oligomer is used as a therapeutic agent for treatment of APC and axin-related disease states.

Assay Systems

The invention provides assay systems and screening methods for identifying specific modulators of LOC169505 activity. As used herein, an "assay system"

encompasses all the components required for performing and analyzing results of an assay that detects and/or measures a particular event. In general, primary assays are used to identify or confirm a modulator's specific biochemical or molecular effect with respect to the LOC169505 nucleic acid or protein. In general, secondary assays further assess the activity of a LOC169505 modulating agent identified by a primary assay and may confirm that the modulating agent affects LOC169505 in a manner relevant to the APC and axin pathways. In some cases, LOC169505 modulators will be directly tested in a secondary assay.

In a preferred embodiment, the screening method comprises contacting a suitable assay system comprising an LOC169505 polypeptide or nucleic acid with a candidate agent under conditions whereby, but for the presence of the agent, the system provides a reference activity (e.g. kinase activity), which is based on the particular molecular event the screening method detects. A statistically significant difference between the agent-biased activity and the reference activity indicates that the candidate agent modulates LOC169505 activity, and hence the APC and axin pathways. The LOC169505 polypeptide or nucleic acid used in the assay may comprise any of the nucleic acids or polypeptides described above.

Primary Assays

The type of modulator tested generally determines the type of primary assay.

Primary assays for small molecule modulators

For small molecule modulators, screening assays are used to identify candidate modulators. Screening assays may be cell-based or may use a cell-free system that recreates or retains the relevant biochemical reaction of the target protein (reviewed in Sittampalam GS *et al.*, Curr Opin Chem Biol (1997) 1:384-91 and accompanying references). As used herein the term "cell-based" refers to assays using live cells, dead cells, or a particular cellular fraction, such as a membrane, endoplasmic reticulum, or mitochondrial fraction. The term "cell free" encompasses assays using substantially purified protein (either endogenous or recombinantly produced), partially purified or crude cellular extracts. Screening assays may detect a variety of molecular events, including protein-DNA interactions, protein-protein interactions (e.g., receptor-ligand binding), transcriptional activity (e.g., using a reporter gene), enzymatic activity (e.g., via a property of the substrate), activity of second messengers, immunogenicity and changes in cellular

morphology or other cellular characteristics. Appropriate screening assays may use a wide range of detection methods including fluorescent, radioactive, colorimetric, spectrophotometric, and amperometric methods, to provide a read-out for the particular molecular event detected.

5 Cell-based screening assays usually require systems for recombinant expression of LOC169505 and any auxiliary proteins demanded by the particular assay. Appropriate methods for generating recombinant proteins produce sufficient quantities of proteins that retain their relevant biological activities and are of sufficient purity to optimize activity and assure assay reproducibility. Yeast two-hybrid and variant screens, and mass
10 spectrometry provide preferred methods for determining protein-protein interactions and elucidation of protein complexes. In certain applications, when LOC169505-interacting proteins are used in screens to identify small molecule modulators, the binding specificity of the interacting protein to the LOC169505 protein may be assayed by various known methods such as substrate processing (e.g. ability of the candidate LOC169505-specific
15 binding agents to function as negative effectors in LOC169505-expressing cells), binding equilibrium constants (usually at least about 10^7 M^{-1} , preferably at least about 10^8 M^{-1} , more preferably at least about 10^9 M^{-1}), and immunogenicity (e.g. ability to elicit LOC169505 specific antibody in a heterologous host such as a mouse, rat, goat or rabbit). For enzymes and receptors, binding may be assayed by, respectively, substrate and ligand
20 processing.

 The screening assay may measure a candidate agent's ability to specifically bind to or modulate activity of a LOC169505 polypeptide, a fusion protein thereof, or to cells or membranes bearing the polypeptide or fusion protein. The LOC169505 polypeptide can be full length or a fragment thereof that retains functional LOC169505 activity. The
25 LOC169505 polypeptide may be fused to another polypeptide, such as a peptide tag for detection or anchoring, or to another tag. The LOC169505 polypeptide is preferably human LOC169505, or is an ortholog or derivative thereof as described above. In a preferred embodiment, the screening assay detects candidate agent-based modulation of LOC169505 interaction with a binding target, such as an endogenous or exogenous protein
30 or other substrate that has LOC169505 -specific binding activity, and can be used to assess normal LOC169505 gene function.

 Suitable assay formats that may be adapted to screen for LOC169505 modulators are known in the art. Preferred screening assays are high throughput or ultra high throughput and thus provide automated, cost-effective means of screening compound

libraries for lead compounds (Fernandes PB, Curr Opin Chem Biol (1998) 2:597-603; Sundberg SA, Curr Opin Biotechnol 2000, 11:47-53). In one preferred embodiment, screening assays uses fluorescence technologies, including fluorescence polarization, time-resolved fluorescence, and fluorescence resonance energy transfer. These systems offer means to monitor protein-protein or DNA-protein interactions in which the intensity of the signal emitted from dye-labeled molecules depends upon their interactions with partner molecules (e.g., Selvin PR, Nat Struct Biol (2000) 7:730-4; Fernandes PB, *supra*; Hertzberg RP and Pope AJ, Curr Opin Chem Biol (2000) 4:445-451).

A variety of suitable assay systems may be used to identify candidate LOC169505 and APC and axin pathways modulators (e.g. U.S. Pat. No. 6,165,992 (kinase assays); U.S. Pat. Nos. 5,550,019 and 6,133,437 (apoptosis assays); and U.S. Pat. Nos. 5,976,782, 6,225,118 and 6,444,434 (angiogenesis assays), among others). Specific preferred assays are described in more detail below.

Protein kinases, key signal transduction proteins that may be either membrane-associated or intracellular, catalyze the transfer of gamma phosphate from adenosine triphosphate (ATP) to a serine, threonine or tyrosine residue in a protein substrate. Radioassays, which monitor the transfer from [gamma-³²P or -³³P]ATP, are frequently used to assay kinase activity. For instance, a scintillation assay for p56 (lck) kinase activity monitors the transfer of the gamma phosphate from [gamma-³³P] ATP to a biotinylated peptide substrate. The substrate is captured on a streptavidin coated bead that transmits the signal (Beveridge M *et al.*, J Biomol Screen (2000) 5:205-212). This assay uses the scintillation proximity assay (SPA), in which only radio-ligand bound to receptors tethered to the surface of an SPA bead are detected by the scintillant immobilized within it, allowing binding to be measured without separation of bound from free ligand. Other assays for protein kinase activity may use antibodies that specifically recognize phosphorylated substrates. For instance, the kinase receptor activation (KIRA) assay measures receptor tyrosine kinase activity by ligand stimulating the intact receptor in cultured cells, then capturing solubilized receptor with specific antibodies and quantifying phosphorylation via phosphotyrosine ELISA (Sadick MD, Dev Biol Stand (1999) 97:121-133). Another example of antibody based assays for protein kinase activity is TRF (time-resolved fluorometry). This method utilizes europium chelate-labeled anti-phosphotyrosine antibodies to detect phosphate transfer to a polymeric substrate coated onto microtiter plate wells. The amount of phosphorylation is then detected using time-

resolved, dissociation-enhanced fluorescence (Braunwalder AF, et al., Anal Biochem 1996 Jul 1;238(2):159-64).

Apoptosis assays. Assays for apoptosis may be performed by terminal
5 deoxynucleotidyl transferase-mediated digoxigenin-11-dUTP nick end labeling (TUNEL)
assay. The TUNEL assay is used to measure nuclear DNA fragmentation characteristic of
apoptosis (Lazebnik *et al.*, 1994, Nature 371, 346), by following the incorporation of
fluorescein-dUTP (Yonehara *et al.*, 1989, J. Exp. Med. 169, 1747). Apoptosis may further
be assayed by acridine orange staining of tissue culture cells (Lucas, R., et al., 1998, Blood
10 15:4730-41). Other cell-based apoptosis assays include the caspase-3/7 assay and the cell
death nucleosome ELISA assay. The caspase 3/7 assay is based on the activation of the
caspase cleavage activity as part of a cascade of events that occur during programmed cell
death in many apoptotic pathways. In the caspase 3/7 assay (commercially available Apo-
ONE™ Homogeneous Caspase-3/7 assay from Promega, cat# 67790), lysis buffer and
15 caspase substrate are mixed and added to cells. The caspase substrate becomes fluorescent
when cleaved by active caspase 3/7. The nucleosome ELISA assay is a general cell death
assay known to those skilled in the art, and available commercially (Roche, Cat#
1774425). This assay is a quantitative sandwich-enzyme-immunoassay which uses
monoclonal antibodies directed against DNA and histones respectively, thus specifically
20 determining amount of mono- and oligonucleosomes in the cytoplasmic fraction of cell
lysates. Mono and oligonucleosomes are enriched in the cytoplasm during apoptosis due
to the fact that DNA fragmentation occurs several hours before the plasma membrane
breaks down, allowing for accumulation in the cytoplasm. Nucleosomes are not present in
the cytoplasmic fraction of cells that are not undergoing apoptosis. An apoptosis assay
25 system may comprise a cell that expresses an LOC169505, and that optionally has
defective APC and axin function (e.g. APC and axin is over-expressed or under-expressed
relative to wild-type cells). A test agent can be added to the apoptosis assay system and
changes in induction of apoptosis relative to controls where no test agent is added, identify
candidate APC and axin modulating agents. In some embodiments of the invention, an
30 apoptosis assay may be used as a secondary assay to test a candidate APC and axin
modulating agents that is initially identified using a cell-free assay system. An apoptosis
assay may also be used to test whether LOC169505 function plays a direct role in
apoptosis. For example, an apoptosis assay may be performed on cells that over- or under-
express LOC169505 relative to wild type cells. Differences in apoptotic response

compared to wild type cells suggests that the LOC169505 plays a direct role in the apoptotic response. Apoptosis assays are described further in US Pat. No. 6,133,437.

Cell proliferation and cell cycle assays. Cell proliferation may be assayed via
5 bromodeoxyuridine (BRDU) incorporation. This assay identifies a cell population undergoing DNA synthesis by incorporation of BRDU into newly-synthesized DNA. Newly-synthesized DNA may then be detected using an anti-BRDU antibody (Hoshino *et al.*, 1986, *Int. J. Cancer* 38, 369; Campana *et al.*, 1988, *J. Immunol. Meth.* 107, 79), or by other means.

10 Cell proliferation is also assayed via phospho-histone H3 staining, which identifies a cell population undergoing mitosis by phosphorylation of histone H3. Phosphorylation of histone H3 at serine 10 is detected using an antibody specific to the phosphorylated form of the serine 10 residue of histone H3. (Chadlee, D.N. 1995, *J. Biol. Chem.* 270:20098-105). Cell Proliferation may also be examined using [³H]-thymidine incorporation (Chen,
15 J., 1996, *Oncogene* 13:1395-403; Jeoung, J., 1995, *J. Biol. Chem.* 270:18367-73). This assay allows for quantitative characterization of S-phase DNA syntheses. In this assay, cells synthesizing DNA will incorporate [³H]-thymidine into newly synthesized DNA. Incorporation can then be measured by standard techniques such as by counting of radioisotope in a scintillation counter (e.g., Beckman LS 3800 Liquid Scintillation
20 Counter). Another proliferation assay uses the dye Alamar Blue (available from Biosource International), which fluoresces when reduced in living cells and provides an indirect measurement of cell number (Voytik-Harbin SL *et al.*, 1998, *In Vitro Cell Dev Biol Anim* 34:239-46). Yet another proliferation assay, the MTS assay, is based on in vitro cytotoxicity assessment of industrial chemicals, and uses the soluble tetrazolium salt,
25 MTS. MTS assays are commercially available, for example, the Promega CellTiter 96[®] Aqueous Non-Radioactive Cell Proliferation Assay (Cat.# G5421).

Cell proliferation may also be assayed by colony formation in soft agar (Sambrook *et al.*, *Molecular Cloning*, Cold Spring Harbor (1989)). For example, cells transformed with LOC169505 are seeded in soft agar plates, and colonies are measured and counted
30 after two weeks incubation.

Cell proliferation may also be assayed by measuring ATP levels as indicator of metabolically active cells. Such assays are commercially available, for example Cell Titer-Glo[™], which is a luminescent homogeneous assay available from Promega.

Involvement of a gene in the cell cycle may be assayed by flow cytometry (Gray JW et al. (1986) Int J Radiat Biol Relat Stud Phys Chem Med 49:237-55). Cells transfected with an LOC169505 may be stained with propidium iodide and evaluated in a flow cytometer (available from Becton Dickinson), which indicates accumulation of cells in different stages of the cell cycle.

Accordingly, a cell proliferation or cell cycle assay system may comprise a cell that expresses an LOC169505, and that optionally has defective APC and axin function (e.g. APC and axin is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the assay system and changes in cell proliferation or cell cycle relative to controls where no test agent is added, identify candidate APC and axin modulating agents. In some embodiments of the invention, the cell proliferation or cell cycle assay may be used as a secondary assay to test a candidate APC and axin modulating agents that is initially identified using another assay system such as a cell-free assay system. A cell proliferation assay may also be used to test whether LOC169505 function plays a direct role in cell proliferation or cell cycle. For example, a cell proliferation or cell cycle assay may be performed on cells that over- or under-express LOC169505 relative to wild type cells. Differences in proliferation or cell cycle compared to wild type cells suggests that the LOC169505 plays a direct role in cell proliferation or cell cycle.

Angiogenesis. Angiogenesis may be assayed using various human endothelial cell systems, such as umbilical vein, coronary artery, or dermal cells. Suitable assays include Alamar Blue based assays (available from Biosource International) to measure proliferation; migration assays using fluorescent molecules, such as the use of Becton Dickinson Falcon HTS FluoroBlock cell culture inserts to measure migration of cells through membranes in presence or absence of angiogenesis enhancer or suppressors; and tubule formation assays based on the formation of tubular structures by endothelial cells on Matrigel® (Becton Dickinson). Accordingly, an angiogenesis assay system may comprise a cell that expresses an LOC169505, and that optionally has defective APC and axin function (e.g. APC and axin is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the angiogenesis assay system and changes in angiogenesis relative to controls where no test agent is added, identify candidate APC and axin modulating agents. In some embodiments of the invention, the angiogenesis assay may be used as a secondary assay to test a candidate APC and axin modulating agents that is initially identified using another assay system. An angiogenesis assay may also be used

to test whether LOC169505 function plays a direct role in cell proliferation. For example, an angiogenesis assay may be performed on cells that over- or under-express LOC169505 relative to wild type cells. Differences in angiogenesis compared to wild type cells suggests that the LOC169505 plays a direct role in angiogenesis. U.S. Pat. Nos.

5 5,976,782, 6,225,118 and 6,444,434, among others, describe various angiogenesis assays.

Hypoxic induction. The alpha subunit of the transcription factor, hypoxia inducible factor-1 (HIF-1), is upregulated in tumor cells following exposure to hypoxia in vitro. Under hypoxic conditions, HIF-1 stimulates the expression of genes known to be
10 important in tumour cell survival, such as those encoding glycolytic enzymes and VEGF. Induction of such genes by hypoxic conditions may be assayed by growing cells transfected with LOC169505 in hypoxic conditions (such as with 0.1% O₂, 5% CO₂, and balance N₂, generated in a Napco 7001 incubator (Precision Scientific)) and normoxic conditions, followed by assessment of gene activity or expression by Taqman®. For
15 example, a hypoxic induction assay system may comprise a cell that expresses an LOC169505, and that optionally has defective APC and axin function (e.g. APC and axin is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the hypoxic induction assay system and changes in hypoxic response relative to controls where no test agent is added, identify candidate APC and axin modulating agents.
20 In some embodiments of the invention, the hypoxic induction assay may be used as a secondary assay to test a candidate APC and axin modulating agents that is initially identified using another assay system. A hypoxic induction assay may also be used to test whether LOC169505 function plays a direct role in the hypoxic response. For example, a hypoxic induction assay may be performed on cells that over- or under-express
25 LOC169505 relative to wild type cells. Differences in hypoxic response compared to wild type cells suggests that the LOC169505 plays a direct role in hypoxic induction.

Cell adhesion. Cell adhesion assays measure adhesion of cells to purified adhesion proteins, or adhesion of cells to each other, in presence or absence of candidate
30 modulating agents. Cell-protein adhesion assays measure the ability of agents to modulate the adhesion of cells to purified proteins. For example, recombinant proteins are produced, diluted to 2.5g/mL in PBS, and used to coat the wells of a microtiter plate. The wells used for negative control are not coated. Coated wells are then washed, blocked with 1% BSA, and washed again. Compounds are diluted to 2× final test concentration

and added to the blocked, coated wells. Cells are then added to the wells, and the unbound cells are washed off. Retained cells are labeled directly on the plate by adding a membrane-permeable fluorescent dye, such as calcein-AM, and the signal is quantified in a fluorescent microplate reader.

5 Cell-cell adhesion assays measure the ability of agents to modulate binding of cell adhesion proteins with their native ligands. These assays use cells that naturally or recombinantly express the adhesion protein of choice. In an exemplary assay, cells expressing the cell adhesion protein are plated in wells of a multiwell plate. Cells expressing the ligand are labeled with a membrane-permeable fluorescent dye, such as
10 BCECF, and allowed to adhere to the monolayers in the presence of candidate agents. Unbound cells are washed off, and bound cells are detected using a fluorescence plate reader.

High-throughput cell adhesion assays have also been described. In one such assay, small molecule ligands and peptides are bound to the surface of microscope slides using a
15 microarray spotter, intact cells are then contacted with the slides, and unbound cells are washed off. In this assay, not only the binding specificity of the peptides and modulators against cell lines are determined, but also the functional cell signaling of attached cells using immunofluorescence techniques in situ on the microchip is measured (Falsey JR et al., Bioconjug Chem. 2001 May-Jun;12(3):346-53).

20

Tubulogenesis. Tubulogenesis assays monitor the ability of cultured cells, generally endothelial cells, to form tubular structures on a matrix substrate, which generally simulates the environment of the extracellular matrix. Exemplary substrates include MatrigelTM (Becton Dickinson), an extract of basement membrane proteins
25 containing laminin, collagen IV, and heparin sulfate proteoglycan, which is liquid at 4°C and forms a solid gel at 37°C. Other suitable matrices comprise extracellular components such as collagen, fibronectin, and/or fibrin. Cells are stimulated with a pro-angiogenic stimulant, and their ability to form tubules is detected by imaging. Tubules can generally be detected after an overnight incubation with stimuli, but longer or shorter time frames
30 may also be used. Tube formation assays are well known in the art (e.g., Jones MK et al., 1999, Nature Medicine 5:1418-1423). These assays have traditionally involved stimulation with serum or with the growth factors FGF or VEGF. Serum represents an undefined source of growth factors. In a preferred embodiment, the assay is performed with cells cultured in serum free medium, in order to control which process or pathway a

candidate agent modulates. Moreover, we have found that different target genes respond differently to stimulation with different pro-angiogenic agents, including inflammatory angiogenic factors such as TNF- α . Thus, in a further preferred embodiment, a tubulogenesis assay system comprises testing an LOC169505's response to a variety of
5 factors, such as FGF, VEGF, phorbol myristate acetate (PMA), TNF- α , ephrin, etc.

Cell Migration. An invasion/migration assay (also called a migration assay) tests the ability of cells to overcome a physical barrier and to migrate towards pro-angiogenic signals. Migration assays are known in the art (e.g., Paik JH et al., 2001, J Biol Chem
10 276:11830-11837). In a typical experimental set-up, cultured endothelial cells are seeded onto a matrix-coated porous lamina, with pore sizes generally smaller than typical cell size. The matrix generally simulates the environment of the extracellular matrix, as described above. The lamina is typically a membrane, such as the transwell polycarbonate membrane (Corning Costar Corporation, Cambridge, MA), and is generally part of an
15 upper chamber that is in fluid contact with a lower chamber containing pro-angiogenic stimuli. Migration is generally assayed after an overnight incubation with stimuli, but longer or shorter time frames may also be used. Migration is assessed as the number of cells that crossed the lamina, and may be detected by staining cells with hemotoxylin solution (VWR Scientific, South San Francisco, CA), or by any other method for
20 determining cell number. In another exemplary set up, cells are fluorescently labeled and migration is detected using fluorescent readings, for instance using the Falcon HTS FluoroBlok (Becton Dickinson). While some migration is observed in the absence of stimulus, migration is greatly increased in response to pro-angiogenic factors. As described above, a preferred assay system for migration/invasion assays comprises testing
25 an LOC169505's response to a variety of pro-angiogenic factors, including tumor angiogenic and inflammatory angiogenic agents, and culturing the cells in serum free medium.

Sprouting assay. A sprouting assay is a three-dimensional *in vitro* angiogenesis
30 assay that uses a cell-number defined spheroid aggregation of endothelial cells ("spheroid"), embedded in a collagen gel-based matrix. The spheroid can serve as a starting point for the sprouting of capillary-like structures by invasion into the extracellular matrix (termed "cell sprouting") and the subsequent formation of complex anastomosing networks (Korff and Augustin, 1999, J Cell Sci 112:3249-58). In an

exemplary experimental set-up, spheroids are prepared by pipetting 400 human umbilical vein endothelial cells into individual wells of a nonadhesive 96-well plates to allow overnight spheroidal aggregation (Korff and Augustin: J Cell Biol 143: 1341-52, 1998). Spheroids are harvested and seeded in 900 μ l of methocel-collagen solution and pipetted
5 into individual wells of a 24 well plate to allow collagen gel polymerization. Test agents are added after 30 min by pipetting 100 μ l of 10-fold concentrated working dilution of the test substances on top of the gel. Plates are incubated at 37°C for 24h. Dishes are fixed at the end of the experimental incubation period by addition of paraformaldehyde. Sprouting intensity of endothelial cells can be quantitated by an automated image analysis system to
10 determine the cumulative sprout length per spheroid.

Primary assays for antibody modulators

For antibody modulators, appropriate primary assays test is a binding assay that tests the antibody's affinity to and specificity for the LOC169505 protein. Methods for
15 testing antibody affinity and specificity are well known in the art (Harlow and Lane, 1988, 1999, *supra*). The enzyme-linked immunosorbant assay (ELISA) is a preferred method for detecting LOC169505-specific antibodies; others include FACS assays, radioimmunoassays, and fluorescent assays.

In some cases, screening assays described for small molecule modulators may also
20 be used to test antibody modulators.

Primary assays for nucleic acid modulators

For nucleic acid modulators, primary assays may test the ability of the nucleic acid modulator to inhibit or enhance LOC169505 gene expression, preferably mRNA
25 expression. In general, expression analysis comprises comparing LOC169505 expression in like populations of cells (*e.g.*, two pools of cells that endogenously or recombinantly express LOC169505) in the presence and absence of the nucleic acid modulator. Methods for analyzing mRNA and protein expression are well known in the art. For instance, Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR (*e.g.*, using
30 the TaqMan®, PE Applied Biosystems), or microarray analysis may be used to confirm that LOC169505 mRNA expression is reduced in cells treated with the nucleic acid modulator (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm DH and Guiseppi-Elie, A Curr

Opin Biotechnol 2001, 12:41-47). Protein expression may also be monitored. Proteins are most commonly detected with specific antibodies or antisera directed against either the LOC169505 protein or specific peptides. A variety of means including Western blotting, ELISA, or in situ detection, are available (Harlow E and Lane D, 1988 and 1999, *supra*).

- 5 In some cases, screening assays described for small molecule modulators, particularly in assay systems that involve LOC169505 mRNA expression, may also be used to test nucleic acid modulators.

Secondary Assays

- 10 Secondary assays may be used to further assess the activity of LOC169505-modulating agent identified by any of the above methods to confirm that the modulating agent affects LOC169505 in a manner relevant to the APC and axin pathways. As used herein, LOC169505-modulating agents encompass candidate clinical compounds or other agents derived from previously identified modulating agent. Secondary assays can also be
15 used to test the activity of a modulating agent on a particular genetic or biochemical pathway or to test the specificity of the modulating agent's interaction with LOC169505.

- Secondary assays generally compare like populations of cells or animals (*e.g.*, two pools of cells or animals that endogenously or recombinantly express LOC169505) in the presence and absence of the candidate modulator. In general, such assays test whether
20 treatment of cells or animals with a candidate LOC169505-modulating agent results in changes in the APC and axin pathways in comparison to untreated (or mock- or placebo-treated) cells or animals. Certain assays use "sensitized genetic backgrounds", which, as used herein, describe cells or animals engineered for altered expression of genes in the APC and axin or interacting pathways.

25

Cell-based assays

- Cell based assays may detect endogenous APC and axin pathways activity or may rely on recombinant expression of APC and axin pathways components. Any of the aforementioned assays may be used in this cell-based format. Candidate modulators are
30 typically added to the cell media but may also be injected into cells or delivered by any other efficacious means.

Animal Assays

A variety of non-human animal models of normal or defective APC and axin pathways may be used to test candidate LOC169505 modulators. Models for defective APC and axin pathways typically use genetically modified animals that have been engineered to mis-express (*e.g.*, over-express or lack expression in) genes involved in the APC and axin pathways. Assays generally require systemic delivery of the candidate modulators, such as by oral administration, injection, etc.

In a preferred embodiment, APC and axin pathways activity is assessed by monitoring neovascularization and angiogenesis. Animal models with defective and normal APC and axin are used to test the candidate modulator's affect on LOC169505 in Matrigel® assays. Matrigel® is an extract of basement membrane proteins, and is composed primarily of laminin, collagen IV, and heparin sulfate proteoglycan. It is provided as a sterile liquid at 4°C, but rapidly forms a solid gel at 37°C. Liquid Matrigel® is mixed with various angiogenic agents, such as bFGF and VEGF, or with human tumor cells which over-express the LOC169505. The mixture is then injected subcutaneously(SC) into female athymic nude mice (Taconic, Germantown, NY) to support an intense vascular response. Mice with Matrigel® pellets may be dosed via oral (PO), intraperitoneal (IP), or intravenous (IV) routes with the candidate modulator. Mice are euthanized 5 - 12 days post-injection, and the Matrigel® pellet is harvested for hemoglobin analysis (Sigma plasma hemoglobin kit). Hemoglobin content of the gel is found to correlate the degree of neovascularization in the gel.

In another preferred embodiment, the effect of the candidate modulator on LOC169505 is assessed via tumorigenicity assays. Tumor xenograft assays are known in the art (see, *e.g.*, Ogawa K et al., 2000, *Oncogene* 19:6043-6052). Xenografts are typically implanted SC into female athymic mice, 6-7 week old, as single cell suspensions either from a pre-existing tumor or from *in vitro* culture. The tumors which express the LOC169505 endogenously are injected in the flank, 1×10^5 to 1×10^7 cells per mouse in a volume of 100 µL using a 27gauge needle. Mice are then ear tagged and tumors are measured twice weekly. Candidate modulator treatment is initiated on the day the mean tumor weight reaches 100 mg. Candidate modulator is delivered IV, SC, IP, or PO by bolus administration. Depending upon the pharmacokinetics of each unique candidate modulator, dosing can be performed multiple times per day. The tumor weight is assessed by measuring perpendicular diameters with a caliper and calculated by multiplying the measurements of diameters in two dimensions. At the end of the experiment, the excised

tumors maybe utilized for biomarker identification or further analyses. For immunohistochemistry staining, xenograft tumors are fixed in 4% paraformaldehyde, 0.1M phosphate, pH 7.2, for 6 hours at 4°C, immersed in 30% sucrose in PBS, and rapidly frozen in isopentane cooled with liquid nitrogen.

5 In another preferred embodiment, tumorigenicity is monitored using a hollow fiber assay, which is described in U.S. Pat No. US 5,698,413. Briefly, the method comprises implanting into a laboratory animal a biocompatible, semi-permeable encapsulation device containing target cells, treating the laboratory animal with a candidate modulating agent, and evaluating the target cells for reaction to the candidate modulator. Implanted cells are
10 generally human cells from a pre-existing tumor or a tumor cell line. After an appropriate period of time, generally around six days, the implanted samples are harvested for evaluation of the candidate modulator. Tumorigenicity and modulator efficacy may be evaluated by assaying the quantity of viable cells present in the macrocapsule, which can be determined by tests known in the art, for example, MTT dye conversion assay, neutral
15 red dye uptake, trypan blue staining, viable cell counts, the number of colonies formed in soft agar, the capacity of the cells to recover and replicate in vitro, etc.

In another preferred embodiment, a tumorigenicity assay use a transgenic animal, usually a mouse, carrying a dominant oncogene or tumor suppressor gene knockout under the control of tissue specific regulatory sequences; these assays are generally referred to as
20 transgenic tumor assays. In a preferred application, tumor development in the transgenic model is well characterized or is controlled. In an exemplary model, the "RIP1-Tag2" transgene, comprising the SV40 large T-antigen oncogene under control of the insulin gene regulatory regions is expressed in pancreatic beta cells and results in islet cell carcinomas (Hanahan D, 1985, Nature 315:115-122; Parangi S et al, 1996, Proc Natl Acad
25 Sci USA 93: 2002-2007; Bergers G et al, 1999, Science 284:808-812). An "angiogenic switch," occurs at approximately five weeks, as normally quiescent capillaries in a subset of hyperproliferative islets become angiogenic. The RIP1-TAG2 mice die by age 14 weeks. Candidate modulators may be administered at a variety of stages, including just prior to the angiogenic switch (e.g., for a model of tumor prevention), during the growth of
30 small tumors (e.g., for a model of intervention), or during the growth of large and/or invasive tumors (e.g., for a model of regression). Tumorigenicity and modulator efficacy can be evaluating life-span extension and/or tumor characteristics, including number of tumors, tumor size, tumor morphology, vessel density, apoptotic index, etc.

Diagnostic and therapeutic uses

Specific LOC169505-modulating agents are useful in a variety of diagnostic and therapeutic applications where disease or disease prognosis is related to defects in the APC and axin pathways, such as angiogenic, apoptotic, or cell proliferation disorders.

- 5 Accordingly, the invention also provides methods for modulating the APC and axin pathways in a cell, preferably a cell pre-determined to have defective or impaired APC and axin function (e.g. due to overexpression, underexpression, or misexpression of APC and axin, or due to gene mutations), comprising the step of administering an agent to the cell that specifically modulates LOC169505 activity. Preferably, the modulating agent
- 10 produces a detectable phenotypic change in the cell indicating that the APC and axin function is restored. The phrase "function is restored", and equivalents, as used herein, means that the desired phenotype is achieved, or is brought closer to normal compared to untreated cells. For example, with restored APC and axin function, cell proliferation and/or progression through cell cycle may normalize, or be brought closer to normal
- 15 relative to untreated cells. The invention also provides methods for treating disorders or disease associated with impaired APC and axin function by administering a therapeutically effective amount of an LOC169505 -modulating agent that modulates the APC and axin pathways. The invention further provides methods for modulating LOC169505 function in a cell, preferably a cell pre-determined to have defective or impaired LOC169505
- 20 function, by administering an LOC169505 -modulating agent. Additionally, the invention provides a method for treating disorders or disease associated with impaired LOC169505 function by administering a therapeutically effective amount of an LOC169505 -modulating agent.

- The discovery that LOC169505 is implicated in APC and axin pathways provides
- 25 for a variety of methods that can be employed for the diagnostic and prognostic evaluation of diseases and disorders involving defects in the APC and axin pathways and for the identification of subjects having a predisposition to such diseases and disorders.

- Various expression analysis methods can be used to diagnose whether LOC169505 expression occurs in a particular sample, including Northern blotting, slot blotting,
- 30 ribonuclease protection, quantitative RT-PCR, and microarray analysis. (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm and Guiseppi-Elie, Curr Opin Biotechnol 2001, 12:41-47). Tissues having a disease or disorder implicating defective APC and axin signaling that

express an LOC169505, are identified as amenable to treatment with an LOC169505 modulating agent. In a preferred application, the APC and axin defective tissue overexpresses an LOC169505 relative to normal tissue. For example, a Northern blot analysis of mRNA from tumor and normal cell lines, or from tumor and matching normal
5 tissue samples from the same patient, using full or partial LOC169505 cDNA sequences as probes, can determine whether particular tumors express or overexpress LOC169505. Alternatively, the TaqMan® is used for quantitative RT-PCR analysis of LOC169505 expression in cell lines, normal tissues and tumor samples (PE Applied Biosystems).

Various other diagnostic methods may be performed, for example, utilizing
10 reagents such as the LOC169505 oligonucleotides, and antibodies directed against an LOC169505, as described above for: (1) the detection of the presence of LOC169505 gene mutations, or the detection of either over- or under-expression of LOC169505 mRNA relative to the non-disorder state; (2) the detection of either an over- or an under-abundance of LOC169505 gene product relative to the non-disorder state; and (3) the
15 detection of perturbations or abnormalities in the signal transduction pathway mediated by LOC169505.

Thus, in a specific embodiment, the invention is drawn to a method for diagnosing a disease or disorder in a patient that is associated with alterations in LOC169505 expression, the method comprising: a) obtaining a biological sample from the patient; b)
20 contacting the sample with a probe for LOC169505 expression; c) comparing results from step (b) with a control; and d) determining whether step (c) indicates a likelihood of the disease or disorder. Preferably, the disease is cancer, most preferably ovarian or pancreatic cancer. The probe may be either DNA or protein, including an antibody.

25 EXAMPLES

The following experimental section and examples are offered by way of illustration and not by way of limitation.

I. C. elegans APC/axin suppressor screen

30 We have discovered that while RNAi of *apr-1* in a wildtype background does not produce a Muv phenotype, *apr-1* inactivation enhances the penetrance of the Muv phenotype of the *pry-1* mutant to 95% (see also Gleason et al., *supra*). This enhancement of the *pry-1* Muv phenotype requires wildtype *bar-1*/beta-catenin and *pop-1*/TCF activity, suggesting that *apr-1* normally negatively regulates beta-catenin. beta-catenin-specific

suppressor genes, when inactivated, likely suppress beta-catenin's inappropriate transcriptional activation of target genes and, therefore, may be relevant for cancer therapy.

- We designed a genetic screen to identify genes in addition to *bar-1*/beta-catenin and *pop-1*/TCF that act positively in beta-catenin signaling and, when inactivated, suppress the Muv mutant phenotype of *pry-1* (*mu38*); *apr-1* (RNAi). The function of individual genes was inactivated by RNAi in *pry-1* mutant L1 larvae, in combination with *apr-1* RNAi, and suppression of the Muv phenotype was scored as a statistically significant increase in the proportion of adults that did not display the Muv phenotype.
- Suppressor genes were subsequently counterscreened to eliminate those that appeared to suppress the *pry-1* (*mu38*); *apr-1* (RNAi) mutant non-specifically, rather than those that specifically function in beta-catenin signaling. Suppressor genes that passed two specificity assays were considered to be beta-catenin-specific suppressors. First, these suppressors, like *bar-1*/beta-catenin, do not suppress the Muv phenotype of three mutations in genes unrelated to beta-catenin signaling (*let-60*/Ras, *lin-12*/Notch, and *lin-15*). Second, these suppressors are not generally defective in the RNAi response, as determined by co-RNAi with genes unrelated to beta-catenin signaling. F26D11.1 was a suppressor of the *muv* phenotype. Orthologs of the modifier are referred to herein as LOC169505.
- BLAST analysis (Altschul et al., *supra*) was employed to identify orthologs of *C.elegans* modifiers. For example, representative sequences from LOC169505, GI# 20537769 (SEQ ID NO:5), shares 41% amino acid identity with the *C.elegans* F26D11.1.
- Various domains, signals, and functional subunits in proteins were analyzed using the PSORT (Nakai K., and Horton P., Trends Biochem Sci, 1999, 24:34-6; Kenta Nakai, Protein sorting signals and prediction of subcellular localization, Adv. Protein Chem. 54, 277-344 (2000)), PFAM (Bateman A., et al., Nucleic Acids Res, 1999, 27:260-2), SMART (Ponting CP, et al., SMART: identification and annotation of domains from signaling and extracellular protein sequences. Nucleic Acids Res. 1999 Jan 1;27(1):229-32), TM-HMM (Erik L.L. Sonnhammer, Gunnar von Heijne, and Anders Krogh: A hidden Markov model for predicting transmembrane helices in protein sequences. In Proc. of Sixth Int. Conf. on Intelligent Systems for Molecular Biology, p 175-182 Ed J. Glasgow, T. Littlejohn, F. Major, R. Lathrop, D. Sankoff, and C. Sensen Menlo Park, CA: AAAI Press, 1998), and clust (Remm M, and Sonnhammer E. Classification of transmembrane protein families in the *Caenorhabditis elegans* genome and identification of human

orthologs. Genome Res. 2000 Nov;10(11):1679-89) programs. For example, the kinase domain of LOC169505 from GI# 20537769 (SEQ ID NO:5) is located at approximately amino acid residues 7 to 186.

5 II. High-Throughput In Vitro Fluorescence Polarization Assay

Fluorescently-labeled LOC169505 peptide/substrate are added to each well of a 96-well microtiter plate, along with a test agent in a test buffer (10 mM HEPES, 10 mM NaCl, 6 mM magnesium chloride, pH 7.6). Changes in fluorescence polarization, determined by using a Fluorolite FPM-2 Fluorescence Polarization Microtiter System
10 (Dynatech Laboratories, Inc), relative to control values indicates the test compound is a candidate modifier of LOC169505 activity.

III. High-Throughput In Vitro Binding Assay.

³³P-labeled LOC169505 peptide is added in an assay buffer (100 mM KCl, 20 mM
15 HEPES pH 7.6, 1 mM MgCl₂, 1% glycerol, 0.5% NP-40, 50 mM beta-mercaptoethanol, 1 mg/ml BSA, cocktail of protease inhibitors) along with a test agent to the wells of a Neutralite-avidin coated assay plate and incubated at 25°C for 1 hour. Biotinylated substrate is then added to each well and incubated for 1 hour. Reactions are stopped by washing with PBS, and counted in a scintillation counter. Test agents that cause a
20 difference in activity relative to control without test agent are identified as candidate APC and axin modulating agents.

IV. Immunoprecipitations and Immunoblotting

For coprecipitation of transfected proteins, 3×10^6 appropriate recombinant cells
25 containing the LOC169505 proteins are plated on 10-cm dishes and transfected on the following day with expression constructs. The total amount of DNA is kept constant in each transfection by adding empty vector. After 24 h, cells are collected, washed once with phosphate-buffered saline and lysed for 20 min on ice in 1 ml of lysis buffer containing 50 mM Hepes, pH 7.9, 250 mM NaCl, 20 mM -glycerophosphate, 1 mM
30 sodium orthovanadate, 5 mM p-nitrophenyl phosphate, 2 mM dithiothreitol, protease inhibitors (complete, Roche Molecular Biochemicals), and 1% Nonidet P-40. Cellular debris is removed by centrifugation twice at $15,000 \times g$ for 15 min. The cell lysate is incubated with 25 μ l of M2 beads (Sigma) for 2 h at 4 °C with gentle rocking.

After extensive washing with lysis buffer, proteins bound to the beads are solubilized by boiling in SDS sample buffer, fractionated by SDS-polyacrylamide gel electrophoresis, transferred to polyvinylidene difluoride membrane and blotted with the indicated antibodies. The reactive bands are visualized with horseradish peroxidase coupled to the appropriate secondary antibodies and the enhanced chemiluminescence (ECL) Western blotting detection system (Amersham Pharmacia Biotech).

V. Kinase assay

A purified or partially purified LOC169505 is diluted in a suitable reaction buffer, e.g., 50 mM Hepes, pH 7.5, containing magnesium chloride or manganese chloride (1-20 mM) and a peptide or polypeptide substrate, such as myelin basic protein or casein (1-10 μ g/ml). The final concentration of the kinase is 1-20 nM. The enzyme reaction is conducted in microtiter plates to facilitate optimization of reaction conditions by increasing assay throughput. A 96-well microtiter plate is employed using a final volume 30-100 μ l. The reaction is initiated by the addition of 33 P-gamma-ATP (0.5 μ Ci/ml) and incubated for 0.5 to 3 hours at room temperature. Negative controls are provided by the addition of EDTA, which chelates the divalent cation (Mg^{2+} or Mn^{2+}) required for enzymatic activity. Following the incubation, the enzyme reaction is quenched using EDTA. Samples of the reaction are transferred to a 96-well glass fiber filter plate (MultiScreen, Millipore). The filters are subsequently washed with phosphate-buffered saline, dilute phosphoric acid (0.5%) or other suitable medium to remove excess radiolabeled ATP. Scintillation cocktail is added to the filter plate and the incorporated radioactivity is quantitated by scintillation counting (Wallac/Perkin Elmer). Activity is defined by the amount of radioactivity detected following subtraction of the negative control reaction value (EDTA quench).

VI. Expression analysis

All cell lines used in the following experiments are NCI (National Cancer Institute) lines, and are available from ATCC (American Type Culture Collection, Manassas, VA 20110-2209). Normal and tumor tissues were obtained from Impath, UC Davis, Clontech, Stratagene, Ardais, Genome Collaborative, and Ambion.

TaqMan[®] analysis was used to assess expression levels of the disclosed genes in various samples.

RNA was extracted from each tissue sample using Qiagen (Valencia, CA) RNeasy kits, following manufacturer's protocols, to a final concentration of 50ng/ μ l. Single stranded cDNA was then synthesized by reverse transcribing the RNA samples using random hexamers and 500ng of total RNA per reaction, following protocol 4304965 of Applied Biosystems (Foster City, CA).

Primers for expression analysis using TaqMan® assay (Applied Biosystems, Foster City, CA) were prepared according to the TaqMan® protocols, and the following criteria: a) primer pairs were designed to span introns to eliminate genomic contamination, and b) each primer pair produced only one product. Expression analysis was performed using a 7900HT instrument.

TaqMan® reactions were carried out following manufacturer's protocols, in 25 μ l total volume for 96-well plates and 10 μ l total volume for 384-well plates, using 300nM primer and 250 nM probe, and approximately 25ng of cDNA. The standard curve for result analysis was prepared using a universal pool of human cDNA samples, which is a mixture of cDNAs from a wide variety of tissues so that the chance that a target will be present in appreciable amounts is good. The raw data were normalized using 18S rRNA (universally expressed in all tissues and cells).

For each expression analysis, tumor tissue samples were compared with matched normal tissues from the same patient. A gene was considered overexpressed in a tumor when the level of expression of the gene was 2 fold or higher in the tumor compared with its matched normal sample. In cases where normal tissue was not available, a universal pool of cDNA samples was used instead. In these cases, a gene was considered overexpressed in a tumor sample when the difference of expression levels between a tumor sample and the average of all normal samples from the same tissue type was greater than 2 times the standard deviation of all normal samples (i.e., $\text{Tumor} - \text{average}(\text{all normal samples}) > 2 \times \text{STDEV}(\text{all normal samples})$).

LOC169505 (SEQ ID NO:1) was overexpressed in 31% of ovarian cancer samples (16 matched sets) and in 38% of pancreas cancer samples (13 matched sets). A modulator identified by an assay described herein can be further validated for therapeutic effect by administration to a tumor in which the gene is overexpressed. A decrease in tumor growth confirms therapeutic utility of the modulator. Prior to treating a patient with the modulator, the likelihood that the patient will respond to treatment can be diagnosed by obtaining a tumor sample from the patient, and assaying for expression of the gene targeted by the modulator. The expression data for the gene(s) can also be used as a

diagnostic marker for disease progression. The assay can be performed by expression analysis as described above, by antibody directed to the gene target, or by any other available detection method.

5 VII. LOC169505 functional assays

RNAi experiments were carried out to knock down expression of LOC169505 (SEQ ID NO:1) in various cell lines using small interfering RNAs (siRNA, Elbashir et al, *supra*).

Effect of LOC169505 RNAi on cell proliferation and growth. BrdU and Cell
10 Titer-Glo™ assays, as described above, were employed to study the effects of decreased LOC169505 expression on cell proliferation. The results of these experiments indicated that RNAi of LOC169505 decreases proliferation in LX1 lung cancer and 231 breast cancer cells.

Effect of LOC169505 RNAi on apoptosis. Nucleosome ELISA apoptosis assay, as
15 described above, was employed to study the effects of decreased LOC169505 expression on apoptosis.

LOC169505 overexpression analysis. LOC169505 (SEQ ID NO:1) was overexpressed and tested in colony growth assays as described above. Overexpressed LOC169505 had moderate morphological effects on cells, and strong effects on colony
20 growth. Effects of overexpressed LOC169505 on expression of various transcription factors was also studied. Overexpressed LOC169505 caused an increased expression of the following transcription factors: SRE (serum response element), AP1 (Activator protein 1), ETS1 (ETS oncogene; v-ets avian erythroblastosis virus e26 oncogene homolog 1), TCF4 (Transcription factor 4; immunoglobulin transcription factor 2) and EGR (Early
25 growth response).

High Throughput Beta Catenin Transcriptional readout assay. This assay is an expanded TaqMan® transcriptional readout assay monitoring changes in the mRNA levels of endogenous beta catenin regulated genes. This assay measures changes in expression of beta catenin regulated cellular genes as a readout for pathway signaling activity.

30 We identified a panel of genes that were transcriptionally regulated by beta catenin signaling, then designed and tested TaqMan® primer/probes sets. We reduced expression of beta catenin by RNAi, and tested its affect on the expression of the transcriptionally regulated genes in multiple cell types. The panel readout was then narrowed to the ten most robust probes.

We then treated cancer cells with siRNAs of the target genes of interest, such as LOC169505, and tested how the reduced levels of the target genes affected the expression levels of the beta catenin regulated gene panel.

Genes that when knocked out via RNAi, demonstrated the same pattern of activity on at least one panel gene as a beta-catenin knockout, were identified as involved in the beta catenin pathway.

TaqMan[®] assays were performed on the RNAs in a 384 well format.

RNAi of LOC169505 showed the same pattern of activity as beta catenin RNAi for at least one of the transcriptionally regulated genes.

High Throughput active nuclear beta catenin measurement assay. Beta catenin is a cytoplasmic gene, which when activated, moves into the nucleus. This assay was designed to measure the amount of active beta catenin protein in the nucleus using an anti active beta catenin antibody and a nuclear staining dye. Using this assay, we looked for genes that when knocked out, decrease beta catenin activity, and hence, the amount of active beta catenin in the nucleus. This assay was performed using Cellomics Inc. instrumentation.

For this assay, cells were transfected in quadruplicate with siRNAs in 96 well format and stained 72 hours post transfection. The amount of nuclear beta catenin was measured using two different methods.

RNAi of LOC169505 caused a decrease in the nuclear beta catenin.

VIII. LOC169505 activity

The purpose of this experiment was identification of LOC169505 activity. Our initial analysis (Example I) had indicated that LOC169505 had enzymatic activity, similar to that of glucokinase. Assays employing various substrates indicated that LOC169505 is in fact a glucokinase, with gluconate as its substrate.